



Fold



Fold

Fold

Fold

Fold

<div><div><div>• First Treatment Dose</div><div>• Step-up Dose 2</div><div>• Step-up Dose 1</div></div><div>Dates of ELREXFIO Injections:</div><div>Healthcare Setting Name:</div><div>After-Hours Phone Number:</div><div>Office Phone Number:</div><div>Name of ELREXFIO Treating Oncologist:</div></div>		<div><div><div><div></div><div></div></div><div><div>IMPORTANT SAFETY INFORMATION FOR PATIENTS RECEIVING TREATMENT WITH ELREXFIO® (elranatamab-bcmm)</div></div></div><div><div>You should always ask your doctor about taking other medications while taking ELREXFIO.</div><div>PP-L1A-USA-0678 © 2025 Pfizer Inc. All rights reserved. December 2025</div><div><div></div><div>Pfizer</div></div></div></div>	
<div><div><div><div></div><div></div></div><div><div>FOR THE PATIENT</div><div>Call your healthcare provider or get emergency help right away if you have any of these symptoms:</div><div><div><div>• Fever 100.4°F (38°C) or higher</div><div>• Trouble breathing</div><div>• Chills</div><div>• Dizziness or light-headedness</div><div>• Fast heartbeat</div><div>• Headache</div><div>• Agitation, trouble staying awake, confusion or disorientation, seeing or hearing things that are not real</div></div><div><div>• Trouble speaking, thinking, remembering things, paying attention, or understanding things</div><div>• Problems walking or muscle weakness</div><div>• Shaking (tremors), loss of balance, or muscle spasms</div><div>• Numbness and tingling (feeling like “pins and needles”)</div><div>• Burning, throbbing, or stabbing pain</div><div>• Changes in your handwriting</div></div></div></div></div></div>		<div><div><div><div></div><div></div></div><div><div>IMPORTANT TO REMEMBER:</div><div>If you have any of these symptoms, call your doctor or seek emergency medical attention right away! These are not all of the possible symptoms of ELREXFIO. Tell your doctor if you have any symptom that bothers you or does not go away.</div></div></div><div><div>You should be hospitalized after administration of each of the first 2 step-up doses within the step-up dosing schedule.</div></div></div>	
<div><div><div><div></div><div></div></div><div><div>FOR HEALTHCARE PROVIDERS</div><div><div><div><div></div><div></div></div><div><div>IMPORTANT SAFETY INFORMATION YOU SHOULD KNOW: ELREXFIO® (elranatamab-bcmm) therapy can cause cytokine release syndrome (CRS) or neurologic toxicity including immune effector cell-associated neurotoxicity syndrome (ICANS), which may be fatal or life threatening. CRS may involve multiple organ systems.</div></div></div></div></div></div></div>		<div><div><div><div></div><div></div></div><div><div>ELREXFIO® REMS Patient Wallet Card</div><div>1-844-923-7845 www.ELREXFIOREMS.com</div></div></div><div><div>Carry this card with you at all times.</div><div><div><div></div><div></div></div><div><div>ELREXFIO® (elranatamab-bcmm)</div><div>INJECTION FOR SUBCUTANEOUS USE 44 mg/1.1 mL 76 mg/1.9 mL</div></div></div><div><div>SHOW THIS CARD</div><div>to any healthcare provider involved in your care and if you go to the emergency room.</div></div></div></div>	

FRONT

REAR